



## REVIEW - CHAPTER 7




## REVIEW - CHAPTER 7



## Integration

- One of the motivating problem for the invention of integral calculus was the calculation of areas and volumes of irregularly shaped regions (quadrature).
- By analytical integration of an integrand function  $f(x)$   
 $\int f(x) dx = F(x) + C$  must be found so that  $F(x)$  is antiderivative of  $f(x) \Rightarrow F'(x) = f(x)$ .
- Using Fundamental theorem of Calculus definite integral will be calculated as:  
 $I(f) = \int_a^b f(x) dx = F(b) - F(a)$
- Some integrals can not be calculated in such a closed form (e.g.,  $f(x) = e^{-x^2}$ ), other are too complicated. So, we often need numerical methods to evaluate numerical integrals.

## Numerical Integration



- $I(f) = \int_a^b f(x) dx = F(b) - F(a) \cong \sum_{i=1}^n (x_{i+1} - x_i) f(\xi_i) = R_n$   
for  $a = x_1 < x_2 < \dots < x_n < x_{n+1} = b$ ,  $\xi_i \in [x_i, x_{i+1}]$ ,  $i=1 \dots n$
- If we have  $\lim_{n \rightarrow \infty} R_n = R$ , where  $R$  is finite, then  $f$  is said to be Riemann Integrable on  $[a, b]$ , and the value of the integral is  $R$ .

## Existence, Conditioning

- **EXISTENCE:**  
The integral exists if the function  $f(x)$  is bounded and has finite number of discontinuities.
- **CONDITIONING:**  
For determination of Conditioning define the infinite norm of  $f(x)$  as:  $\text{norm}_\infty(f) = \max_{x \in [a, b]} |f(x)|$ .  
Suppose that  $f^\wedge$  is a perturbation of the integrand function  $f$ , then we have:  
 $\Delta(\text{output}) = |I(f^\wedge) - I(f)| = \left| \int_a^b f^\wedge(x) dx - \int_a^b f(x) dx \right| = \int_a^b |f^\wedge(x) - f(x)| dx = (b-a) \text{norm}_\infty(f^\wedge - f) = \text{cond} * \Delta(\text{input})$ .  
Cond is at most  $(b-a) \rightarrow$  well conditioned (smoothing).

## Quadrature Rule Derived by Polynomials

- For  $n$  points the interpolating polynomial of order  $(n-1)$  can be derived (e.g., represented in the Lagrange form):  
 $p_{n-1}(x) = f(x_1) l_1(x) + \dots + f(x_n) l_n(x)$
- Integral of interpolant is taken as an approximate integral of the original function  $f(x)$ :  
 $\int_a^b p_{n-1}(x) dx = f(x_1) \int_a^b l_1(x) dx + \dots + f(x_n) \int_a^b l_n(x) dx$   
The above expression is the quadrature rule with weights equal to integrals of Lagrange basis functions.

## Accuracy Estimation

- $|I(f) - Q_n(f)| = |I(f) - I(p_{n-1})|$   
 (from the previous slide on the quadrature approximation)  
 $\leq (b-a) \text{norm}_\infty(f - p_{n-1})$   
 (from the error of interpolation - Chap 5)  
 $\leq (1/4n) (b-a) h^n \text{norm}_\infty(f^{(n)})$   
 (because  $(b-a) = n \cdot h$ )  
 $\leq (1/4) h^{n+1} \text{norm}_\infty(f^{(n)})$ .
- With larger number of points  $n$  or/and smaller  $h$  the accuracy can be improved.

## Stability of Quadrature Rules

- $|Q_n(f^{\wedge}) - Q_n(f)| = |Q_n(f^{\wedge} - f)| = |\sum_{i=1}^n w_i (f^{\wedge}(x_i) - f(x_i))|$   
 $\leq \sum_{i=1}^n (|w_i| * |f^{\wedge}(x_i) - f(x_i)|)$   
 $\leq (\sum_{i=1}^n |w_i|) \text{norm}_\infty(f^{\wedge} - f)$ .
- From the first moment equation we know that  $\sum_{i=1}^n |w_i| = (b-a)$ .
- If all weights are nonnegative then the absolute condition number of the quadrature rule is  $(b-a)$  (same as integration) - stable quadrature rule.
- If some weights are negative then the absolute condition number of the quadrature rule can be much larger - unstable quadrature rule.

## Change of Interval $[\alpha, \beta] \rightarrow [a, b]$

- $\int_a^b f(x) dx \Rightarrow \int_a^b g(t) dt = I(g)$   
 $t = [(b-a)x + a\beta - b\alpha] / (\beta - \alpha) \quad dt/dx = (b-a) / (\beta - \alpha)$
- $\int_a^b g(t) dt = (b-a) / (\beta - \alpha) \int_a^b g([(b-a)x + a\beta - b\alpha] / (\beta - \alpha)) dx$   
 $\approx (b-a) / (\beta - \alpha) \sum_{i=1}^n w_i g([(b-a)x_i + a\beta - b\alpha] / (\beta - \alpha))$
- Example: Use two-point Gaussian quadrature  $G_2$  on  $[-1, 1]$  to approximate the integral:  $I(g) = \int_0^1 e^{-t^2} dt$
- $x_1 = -1/\sqrt{3}, x_2 = 1/\sqrt{3}, w_1 = w_2 = 1,$   
 $t = (x+1)/2, (b-a) / (\beta - \alpha) = 1/2$   
 $G_2(g) = 1/2 [ e^{-((x_1+1)/2)^2} + e^{-((x_2+1)/2)^2} ] \approx 0.746595$   
 which is more accurate as by Simpson's rule, despite using only two points.

## Adaptive quadrature

```

procedure adaptquad(f, a, b, I^)
  I1=Qn1(f, a, b) //evaluate quadrature rules
  I2=Qn2(f, a, b)
  m=a+(b-a)/2 //compute midpoint of the
  interval
  if (m<=a) or (m>=b) //if no more machine numbers
    warning //tolerance may not be met
    return I2 //return best result
  end
  if I^+(I2-I1) = I^ then //converg. tolerance met
    return I2 //return converged result
  else //refine recursively
    return(adaptquad(f, a, m, I^) +
           adaptquad(f, m, b, I^))
  end
end
    
```

## Integral Equation

- Integral equations arise in observational sciences (e.g., astronomy, seismology, spectrometry). A typical representative of integral equations is the Fredholm IE of the first kind
 
$$\int_a^b K(s, t) u(t) dt = f(s)$$
 $K$  and  $f$  are known,  $u$  must be determined.
- We have to determine a function, which results in known definite integral. It is obvious that this problem could be ill-conditioned, because many different functions can have same integrals.
- Solution procedure:
  - discretize  $s$ , and  $t$ ,
  - Replace the integral by quadrature rule,
  - Solve the resulting linear system.

## Integral Equation

- IEs arise in observational sciences (e.g., astronomy, seismology, spectrometry).  $K$  - response function of an calibrated instrument,  $f$  - measured data and,  $u$  - signal to be sought.
- In effect, we try to resolve the measured data as a linear combination of standard signals.
- $\int_{-1}^1 (1 - a^*s^*t) u(t) dt = 1, K = 1 - a^*s^*t, f(s) = 1, a > 0$ 
  - taking 2 subintervals and using midpoint rule:  $t_1 = -1/2, t_2 = 1/2, w_1 = w_2 = 1$ . Taking also  $s_1 = -1/2, s_2 = 1/2$  we get:
 
$$Ax = [1+a/4, 1-a/4; 1-a/4, 1+a/4]^* [x_1; x_2] = [1; 1] = y$$
 with the solution  $x = [1/2; 1/2]$ , which is independent of  $a$ .
- If  $a$  is small (insensitive instrument),  $A$  becomes near singular, ill-conditioned, error can be arbitrary large.